

# AmeriStand 518NT



#### FALL DORMANCY: 5.2

#### WINTERHARDINESS: 1.9

#### **CUTTINGS/SEASON: 4-5**

#### PERFORMANCE PROFILE

Traffic Tested®: Excellent
Yield Potential: Excellent
Stand Persistence: Excellent
Recovery After Cutting: V. Fast

#### **RESISTANCE RATINGS**

Phytophthora Root Rot: HR

Aphanomyces Root Rot RACE 1: HR

Anthracnose: HR

Verticillium Wilt: HR

Bacterial Wilt: HR

Fusarium Wilt: HR

Pea Aphid: HR

Stem Nematode: HR

Root Knot Nematode: HR

HR = >51% Resistance R = 31-50% Resistance MR = 15-30% Resistance LR = 6-14% Resistance

# **High Yield Potential with Nematode Resistance**

- Widely adapted variety with excellent winter hardiness
- High resistance to nine yield-robbing diseases and pests
- Very fast recovery for frequent harvest schedules under intensive management



## NEMATODES ARE YIELD ROBBERS

**Stem Nematodes** infestations can cause stunted plants and thin stands. Under warm, humid conditions, they can migrate into leaf tissue, killing chloroplasts and turning the leaves white. Infestations can cause stunted plants and thin stands. Under warm, humid conditions, they can migrate into leaf tissue, killing chloroplasts and turning the leaves white.

**Root Knot Nematodes** are among the most widespread and economically damaging to alfalfa. They are most abundant in sandy loam soils and infect roots, causing galls and lateral root growth. Bacterial wilt, *Phytophthora* root rot, *Fusarium* wilt and stem nematode damage may be enhanced when Northern root knot is present.



## VARIETY PERFORMANCE

YIELD TRIAL LOCATION	TRIAL YEARS REPORTED	MULTI-YEAR TOTAL TONS PER ACRE	MULTI-YEAR % OF MEAN
Fruita, CO	3	17.88	106%
Rexburg, ID	5	29.51	112%
Nampa, ID	3	30.36	111%
Touchet, WA	4	44.67	108%
Othello, WA	2	24.58	107%
Butter Creek, OR	5	43.89	103%

The above table compares variety performance in locations with positive results relative to trial means.

America's Alfalfa" and Traffic Tested" are registered trademarks of Forage Genetics International, LLC. ©2017 Forage Genetics International, LLC. Due to factors outside of Forage Genetics International's (FBI's) control, such as weather, crop production patterns, and other factors, results to be obtained, including but not limited to yields or financial performance, cannot be predicted or guaranteed by FGI. Results are based upon FGI controlled tests and field trials and public trials. Actual results may vary.

